# The Influence of Selected Risk Factors on Gestational Diabetes Across Ethnic/Racial Subgroups in Utah



## Introclustion

Figure 1. Trends in Gestational Diabetes

among Utah Mothers 1997-2001

- Nationally, gestational diabetes affects about 4 percent of all pregnancies
- Gestational diabetes in Utah is lower than the U.S., but has increased

consistently in the past five years to the current rate of 2.3% (Figure 1

- Risks for infants born to mothers with gestational diabetes include:
- Increased risk of shoulder dystocia
- Increased risk of macrosomia
- Increased risk of birth injuries
- Increased risk of hypoglycemia
- Risks for mothers with gestational diabetes include:

Source: Office of Vital Records and Statistics

- Increased risk of developing type 2 diabetes
- Increased risk of permanent change in lipid metabolism after pregnancy



### Methods

### Research Questions


- Are there differences in gestational diabetes across racial/ethnic groups that persist with known risk factors controlled?
- Do these risk factors operate differently across racial/ethnic groups?

### Methodology

- Data are from Utah Office of Vital Records and Statistics
- Study was limited to singleton births for non-Hispanic/Latina white, Hispanic/Latina and Native American mothers (Figure 2)
- Births to mothers with pre-existing diabetes were excluded from the analysis
- Logistic regression was used to assess the risk of gestational diabetes
- Controls included age, pre-pregnancy weight status, number of prior live

births, and number of prior still births

- Analyses were run for the entire population; race and ethnicity were included in the model
- Analyses were also run for each ethnic/racial subgroup





## Findings

**Mothers by Race/Ethnicity, 1999-2001** 

 Less than 2% (1.9%) of non-Hispanic/Latina white mothers had gestational diabetes, compared to 2.8% of Hispanic/Latina mothers, and 5.4% of Native

American mothers Figure 3. Prevalence of Gestational Diabetes for Utah (Figure 3)

For each racial or ethnic group, with all known risk factors controlled:

Age and obesity increased the risk of gestational diabetes for all

**Race or Ethnicity** Number of prior

Source: Office of Vital Records and Statistics live births and prior stillbirths did not increase the risk of gestational diabetes for any group

- Unlike obesity, being moderately overweight does not appear to be a significant risk factor for Native American mothers
- Hispanic/Latina mothers were 1.5 times more likely to have gestational diabetes than non-Hispanic/Latina white mothers (with controls)
- Native American mothers were 2.2 times more likely to have gestational diabetes than non-Hispanic/Latina white mothers (with controls)



# Implications

- The higher prevalence of gestational diabetes seen in Hispanic/Latina and Native American mothers persists even with known risk factors controlled
- This finding suggests that other unmeasured factors influence the development of gestational diabetes
- We may need to pay closer attention to the future risk of heart disease and type 2 diabetes for women who have had gestational diabetes
- We may need to make special efforts to make sure our Hispanic/Latina and Native American mothers do not fall out of the medical loop after delivery

#### References

Oostlund, I Hanson U, Bjoorklunk A, Hjhererg R, Eva N, Nordlander, E, Marja-Liisa, S Wager J. 2003. Maternal and Fetal Outcomes if Gestational Impaired Glucose Tolerance is Not Treated. Diabetes Care 26: 2107-2111

Utah Office of Vital Records and Statistics 1990-2001, Utah Department of Health, Salt Lake City, UT

Xiong, X, Saunders I d, Want F L, Demianczuk, NN. 2001. Gestational Diabetes Mellitus: Prevalence, Risk Factors, Maternal and Infant Outcomes. International Journal of Gynecology and Obstetrics 75:221-228.

#### Presented by

Brenda Ralls, PhD William F. Stinner, PhD Shaheen Hossain,PhD Nancy C. Neff, MS LaDene Larsen, RN Terry Allen, PhD Jeffrey Duncan, MS Nathan Bardeen, BS Karen Zinner, MPH

